

Register [here](#) for access to Compass SHARP Resources

Section 1: DNV Standards

DNV Standard	Compass SHARP Recommendation	Compass SHARP Resources to Meet Standard	Supporting Evidence & Other Guidelines
DNV NIAHO Standard PE.3 SR.1 – Hospitals must implement safe opioid prescribing practices	Adopt procedure-specific opioid prescribing limits and defaults within the EHR; implement discharge education for all surgical patients.	Opioid Medication Dictionary ; Guidelines on Limiting Opioids in Perioperative Settings ; Compass Opioid Prescribing + Treatment Guidance Toolkit Education Toolkit .	Michigan OPEN prescribing recommendations ; ERAS Society multimodal analgesia protocols ; CDC Opioid Prescribing Guideline (2022) ; Comagine Health Opioid Stewardship Guide .
DNV NIAHO Standard PC.3 SR.5 – Hospitals must provide naloxone and overdose prevention education to at-risk patients.	Ensure naloxone is prescribed and dispensed at discharge for patients with chronic opioid use, OUD, or high-dose prescriptions.	Naloxone Clinical Toolkit ; Staff Training Module on Patient Education .	CDC Naloxone Guidance ; SAMHSA TIP 63 ; CMS Opioid Roadmap (Naloxone provision) .

Section 2: Joint Commission Standards

Joint Commission Standard	Compass SHARP Recommendation	Compass SHARP Resources to Meet Standard	Supporting Evidence & Other Guidelines
NPSG.03.05.01 – Reduce likelihood of patient harm associated with opioid use.	Standardized opioid risk screening in pre-op and admission workflows; implement opioid stewardship committee to review outcomes.	<u>Compass SHARP Risk Screening Tool</u> ; <u>Compass SHARP 5-Step Implementation Tool</u>	<u>Joint Commission R3 Report on Opioid Stewardship (2018)</u> ; <u>CDC Guideline (2022)</u> ; <u>ERAS protocols</u> .
PC.01.02.07 – Hospitals must manage patients with addiction and provide continuity of care.	Continue MOUD (buprenorphine, methadone) perioperatively; ensure addiction consult services are available.	<u>Perioperative Management of Patients on Opioids (Including MOUD)</u>	<u>SAMHSA TIP 63</u> ; <u>Hah et al., 2017 (Anesthesia & Analgesia)</u> ; <u>Comagine recommendations</u> .

Section 3: CMS Measures & Conditions of Participation

CMS Measure / Condition	Compass SHARP Recommendation	Compass SHARP Resources to Meet Standard	Supporting Evidence & Other Guidelines
CMS Opioid Misuse Strategy – Safe prescribing and monitoring.	Implement audit & feedback on surgical opioid prescribing; track opioid-naïve → chronic use rates.	<u>PDSA Cycle Samples</u> .	<u>CMS Opioid Roadmap (2019)</u> ; <u>Michigan OPEN outcomes data</u> ; <u>Oderda et al., 2020 (Quality indicators)</u> .
CMS Quality Reporting Program – Opioid-related adverse event reduction.	Standardize naloxone distribution program and patient/family education.	<u>Naloxone Distribution Protocol</u> ; <u>Patient Education Handouts</u> .	<u>CDC Naloxone Guidance (2024)</u> ; <u>Joint Commission R3 Report</u> ; <u>State-level naloxone laws</u> .