



Non-Pharmacologic Treatments

Although nonopioid medications are recommended as first-line *pharmacologic* treatment for pain, non-pharmacological therapy is consistently regarded as the preferred initial modality of treatment. Below are descriptions of some of these therapies and associated supporting evidence.

Lifestyle Modification

Exercise and sleep restoration are both supported in the literature as effective interventions to modify and/or prevent pain. Exercise has specifically been linked to both nociceptive (such as chronic lower back) and central sensitization (such as fibromyalgia) type pain improvement, and is a relatively easy, self-directed therapy to recommend for capable patients.^{1,2} Sleep restoration, while straight-forward in some patients, can be more complicated in others. Studies suggest that cognitive behavioral therapy can be an important piece of both improving sleep quality, which in turn is linked to chronic pain development, as well as decreasing pain severity.^{3,4}

Physiotherapy Options

Active therapies such as physical and occupational therapy, therapeutic exercise, aquatic therapy, self-directed activity, neuromuscular re-education, activities of daily living, and functional activities are widely used and accepted methods of care for a variety of types of pain. There is moderate to strong evidence proving the efficacy of exercise therapy for pain relief and functional improvement in patients with musculoskeletal pain.⁵ Massage is the manipulation of soft tissue and may include stimulation of acupuncture points and acupuncture channels (acupressure), application of suction cups, and techniques that include pressing, lifting, rubbing, pinching of soft tissues by or with the practitioner's hands. There is good evidence that massage therapy in combination with exercise reduces pain and improves function in the short term for patients with subacute low back pain.⁶⁻⁸ There is some evidence that 10 weeks of either relaxation massage or structural massage are more effective than usual care and equally effective in improving functional disability and reducing symptoms of pain in people with chronic low back pain with benefits lasting at least six months.⁹ There is also some evidence that in the setting of chronic neck pain four weeks of weekly hour-long massage leads to benefits in both pain and function, and there are incremental benefits from multiple massage sessions per week (up to three sessions) over a single massage session.¹⁰

Psychotherapeutic Interventions

Psychotherapeutic interventions are recommended as a component of treatment for patients with chronic pain; they may also have utility for some patients with acute pain. Cognitive and behavioral interventions have a neurophysiological basis and are well established as diagnostic and therapeutic modalities.¹¹⁻¹³ Patients without behavioral health diagnoses may benefit from interventions that aid in developing better strategies to cope with pain or adjust to disability. A psychologist with a PhD, PsyD, or



Compass Opioid Prescribing + Treatment Guidance Toolkit

EdD or psychiatric MD/DO can perform psychosocial treatment through individual or group therapy. For chronic pain, other licensed mental health providers, licensed health care providers with training in CBT or certified CBT therapists who have experience in treating chronic pain disorders in injured workers may perform treatment in consultation with a PhD, PsyD, EdD or psychiatric MD/DO.

Complementary and Alternative Treatments

Manipulation encompasses a variety of modalities including osteopathic manipulative treatment, chiropractic manipulative treatment, manual therapy, manipulation or mobilization. There is good evidence that manipulation can facilitate pain reduction and improved function for both spinal and extremity injuries.¹⁴⁻¹⁸ Acupuncture is the insertion and removal of filiform needles to stimulate acupoints (acupuncture points) and is recommended for subacute or chronic pain (including low back and knee pain) patients who are trying to increase function, decrease medication usage and have an expressed interest in this modality. There is evidence supporting acupuncture use in reduction of disability and pain in chronic low back pain patients and there is some evidence that acupuncture is better than no acupuncture for axial chronic low back pain.¹⁹⁻²² There is also evidence supporting acupuncture for reduction of pain or improvement of function among patients older than 50 years with moderate to severe chronic knee pain from symptoms of osteoarthritis.²³ If not otherwise within their professional scope of practice and licensure, those performing acupuncture should have the appropriate credentials, such as LAc, RAc or DiplAc. Other phyto-chemicals and dietary supplements for various pain conditions that have evidence for short-term use include, but are not limited to, avocado-soybean unsaponifiables²⁴⁻²⁶, collagen hydrolysate²⁷⁻²⁹, passion fruit peel extract³⁰, Curcuma longa extract³¹⁻³⁴, Boswellia serrata extract³⁵⁻³⁷, curcumin³⁸⁻⁴⁰, pycnogenol^{41,42}, L-carnitine^{43,44}, undenatured type II collagen⁴⁵, methylsulfonylmethane⁴⁶, diacerein⁴⁷, glucosamine^{46,48,49}, chondroitin⁵⁰, capsaicin^{51,52}, alpha-lipoic acid^{53,54}, and thiamine⁵⁵.

Procedure-Based Interventions

Trigger point injections involve injection of a corticosteroid/anesthetic/saline combination into a tensed muscle. Indications supported by evidence include a palpable taut band or nodule, reproducible pain with palpation and/or a chronic painful conditions.⁵⁶⁻⁵⁹ Trigger point injections also have been found to be a successful treatment strategy for migraines.^{60,61} Trigger point injections of corticosteroids are most effective for adhesive capsulitis, rotator cuff tendinopathy, impingement syndrome and tendon disorders. Trigger point dry needling is a skilled intervention that utilizes a solid filament needle to penetrate the skin and underlying tissues to treat muscular, neural and other connective tissues for the evaluation and management of neuro-musculoskeletal conditions, pain, movement impairments and disability. A 2017 systematic review of 15 studies suggests that dry needling is effective in the short term for pain relief, increases range of motion and improves quality of life when compared to no intervention, sham or placebo.⁶² TENS treats pain by delivering small electrical impulses through electrodes that flood pain receptors in the body, reducing their ability to transmit pain signals to the brain. Some good-quality systematic reviews suggest that TENS is effective for musculoskeletal and postoperative pain.⁶³⁻⁶⁶



Compass Opioid Prescribing + Treatment Guidance Toolkit

Practical Application

The above interventions can be split into two categories, self-directed after training vs professionally directed:

Self-Directed After Training	Professionally Directed
Structured exercise	Physical approach
Mind-body therapies	+ Acupuncture
+ Biofeedback	+ Assistive devices
+ Movement meditation	+ Blocks/ablation
+ Mindfulness	+ Ergonomic modifications
Relaxation	+ Light therapy
Music	+ Massage
Neurostimulators	+ Osteo-manipulation
Nutritional approach	+ Physical therapy/occupational therapy
Thermal modalities/balneotherapy	+ Regenerative therapies (platelet rich plasma, prolotherapy, stem cells)
	+ Stimulators (peripheral, spinal, deep brain)
	+ Surgeries
	+ Trigger point interventions
	+ Ultrasound
Sleep hygiene	Psychological approaches/pain behavior therapies
Spiritual practices	+ Acceptance and commitment therapy
Tobacco cessation	+ Cognitive behavioral therapy/ behavioral therapies
Weight reduction	+ Pain education

References

1. Mior S. Exercise in the Treatment of Chronic Pain. The Clinical Journal of Pain. 2001;17(4):S77.
2. Landmark T, Romundstad P, Borchgrevink PC, Kaasa S, Dale O. Associations between recreational exercise and chronic pain in the general population: Evidence from the HUNT 3 study. PAIN®. 2011;152(10):2241-2247. doi:10.1016/j.pain.2011.04.029
3. Smith MT, Haythornthwaite JA. How do sleep disturbance and chronic pain inter-relate? Insights from the longitudinal and cognitive-behavioral clinical trials literature. Sleep Medicine Reviews. 2004;8(2):119-132. doi:10.1016/S1087-0792(03)00044-3
4. Tang NKY, Goodchild CE, Webster LR. Sleep and Chronic Pain. In: Deer TR, Leong MS, Ray AL, eds. Treatment of Chronic Pain by Integrative Approaches: The AMERICAN ACADEMY of PAIN MEDICINE Textbook on Patient Management. Springer; 2015:203-217. doi:10.1007/978-1-4939-1821-8_16
5. Babatunde OO, Jordan JL, Van der Windt DA, Hill JC, Foster NE, Protheroe J. Effective treatment options for musculoskeletal pain in primary care: A systematic overview of current evidence. PLoS One. 2017;12(6). doi:10.1371/journal.pone.0178621
6. Cherkin DC, Eisenberg D, Sherman KJ, et al. Randomized Trial Comparing Traditional Chinese



Compass Opioid Prescribing + Treatment Guidance Toolkit

- Medical Acupuncture, Therapeutic Massage, and Self-care Education for Chronic Low Back Pain. *Arch Intern Med.* 2001;161(8):1081-1088. doi:10.1001/archinte.161.8.1081
7. Furlan AD, Imamura M, Dryden T, Irvin E. Massage for low-back pain. *Cochrane Database Syst Rev.* 2008;(4):CD001929. doi:10.1002/14651858.CD001929.pub2
 8. Preyde M. Effectiveness of massage therapy for subacute low-back pain: a randomized controlled trial. *CMAJ.* 2000;162(13):1815-1820.
 9. Cherkin DC, Sherman KJ, Kahn J, et al. A Comparison of the Effects of 2 Types of Massage and Usual Care on Chronic Low Back Pain: A Randomized, Controlled Trial. *Ann Intern Med.* 2011;155(1):1-9. doi:10.1059/0003-4819-155-1-201107050-00002
 10. Sherman KJ, Cook AJ, Wellman RD, et al. Five-Week Outcomes From a Dosing Trial of Therapeutic Massage for Chronic Neck Pain. *Ann Fam Med.* 2014;12(2):112-120. doi:10.1370/afm.1602
 11. Jensen MP. A Neuropsychological Model of Pain: Research and Clinical Implications. *The Journal of Pain.* 2010;11(1):2-12. doi:10.1016/j.jpain.2009.05.001
 12. Wetherell JL, Afari N, Rutledge T, et al. A randomized, controlled trial of acceptance and commitment therapy and cognitive-behavioral therapy for chronic pain. *PAIN.* 2011;152(9):2098-2107. doi:10.1016/j.pain.2011.05.016
 13. Ehde DM, Dillworth TM, Turner JA. Cognitive-behavioral therapy for individuals with chronic pain: efficacy, innovations, and directions for research. *Am Psychol.* 2014;69(2):153-166. doi:10.1037/a0035747
 14. Rubinstein SM, Zoete A de, Middelkoop M van, Assendelft WJJ, Boer MR de, Tulder MW van. Benefits and harms of spinal manipulative therapy for the treatment of chronic low back pain: systematic review and meta-analysis of randomised controlled trials. *BMJ.* 2019;364. doi:10.1136/bmj.l689
 15. Chou R, Deyo R, Friedly J, et al. Noninvasive Treatments for Low Back Pain. Agency for Healthcare Research and Quality (US); 2016. Accessed March 19, 2020. <http://www.ncbi.nlm.nih.gov/books/NBK350276/>
 16. Gross A, Langevin P, Burnie SJ, et al. Manipulation and mobilisation for neck pain contrasted against an inactive control or another active treatment. *Cochrane Database of Systematic Reviews.* 2015;(9). doi:10.1002/14651858.CD004249.pub4
 17. Rubinstein SM, Middelkoop M van, Assendelft WJ, Boer MR de, Tulder MW van. Spinal manipulative therapy for chronic low-back pain. *Cochrane Database of Systematic Reviews.* 2011;(2). doi:10.1002/14651858.CD008112.pub2
 18. Jansen MJ, Viechtbauer W, Lenssen AF, Hendriks EJM, de Bie RA. Strength training alone, exercise therapy alone, and exercise therapy with passive manual mobilisation each reduce pain and disability in people with knee osteoarthritis: a systematic review. *Journal of Physiotherapy.* 2011;57(1):11-20. doi:10.1016/S1836-9553(11)70002-9
 19. Chou R, Gordon DB, de Leon-Casasola OA, et al. Management of Postoperative Pain: A Clinical Practice Guideline From the American Pain Society, the American Society of Regional Anesthesia and Pain Medicine, and the American Society of Anesthesiologists' Committee on Regional Anesthesia, Executive Committee, and Administrative Council. *The Journal of Pain.* 2016;17(2):131-



Compass Opioid Prescribing + Treatment Guidance Toolkit

157. doi:10.1016/j.jpain.2015.12.008
20. Haake M, Müller H-H, Schade-Brittinger C, et al. German Acupuncture Trials (Gerac) For Chronic Low Back Pain: Randomized, Multicenter, Blinded, Parallel-Group Trial With 3 Groups. *Arch Intern Med.* 2007;167(17):1892-1898. doi:10.1001/Archinte.167.17.1892
21. Cherkin DC, Sherman KJ, Avins AL, et al. A Randomized Trial Comparing Acupuncture, Simulated Acupuncture, and Usual Care for Chronic Low Back Pain. *Arch Intern Med.* 2009;169(9):858-866. doi:10.1001/archinternmed.2009.65
22. Brinkhaus B, Witt CM, Jena S, et al. Acupuncture in Patients With Chronic Low Back Pain: A Randomized Controlled Trial. *Arch Intern Med.* 2006;166(4):450-457. doi:10.1001/archinte.166.4.450
23. Hinman RS, McCrory P, Pirotta M, et al. Acupuncture for Chronic Knee Pain: A Randomized Clinical Trial. *JAMA.* 2014;312(13):1313-1322. doi:10.1001/jama.2014.12660
24. Thierry Appelboom J-YR Joseph Schuermans, Gust Verbruggen, Yves Henrotin. Symptoms modifying effect of avocado/soybean unsaponifiables (ASU) in knee osteoarthritis. *Scandinavian Journal of Rheumatology.* 2001;30(4):242-247. doi:10.1080/030097401316909602
25. Christiansen BA, Bhatti S, Goudarzi R, Emami S. Management of Osteoarthritis with Avocado/Soybean Unsaponifiables. *CARTILAGE.* 2015;6(1):30-44. doi:10.1177/1947603514554992
26. Salehi B, Rescigno A, Dettori T, et al. Avocado–Soybean Unsaponifiables: A Panoply of Potentialities to Be Exploited. *Biomolecules.* 2020;10(1):130. doi:10.3390/biom10010130
27. Moskowitz RW. Role of collagen hydrolysate in bone and joint disease. *Seminars in Arthritis and Rheumatism.* 2000;30(2):87-99. doi:10.1053/sarh.2000.9622
28. Bello AE, Oesser S. Collagen hydrolysate for the treatment of osteoarthritis and other joint disorders: a review of the literature. *Current Medical Research and Opinion.* 2006;22(11):2221-2232. doi:10.1185/030079906X148373
29. Song H, Li B. Beneficial Effects of Collagen Hydrolysate: A Review on Recent Developments. *Biomed J Sci & Tech Res.* 2017;1:1-4.
30. Farid R, Rezaieyazdi Z, Mirfeizi Z, et al. Oral intake of purple passion fruit peel extract reduces pain and stiffness and improves physical function in adult patients with knee osteoarthritis. *Nutrition Research.* 2010;30(9):601-606. doi:10.1016/j.nutres.2010.08.010
31. Madhu K, Chanda K, Saji MJ. Safety and efficacy of Curcuma longa extract in the treatment of painful knee osteoarthritis: a randomized placebo-controlled trial. *Inflammopharmacol.* 2013;21(2):129-136. doi:10.1007/s10787-012-0163-3
32. Bethapudi B, Murugan S, Illuri R, Mundkinajeddu D, Velusami CC. Bioactive Turmerosaccharides from Curcuma longa Extract (NR-INF-02): Potential Ameliorating Effect on Osteoarthritis Pain. *Pharmacogn Mag.* 2017;13(Suppl 3):S623-S627. doi:10.4103/pm.pm_465_16
33. Henrotin Y, Malaise M, Wittoek R, et al. Bio-optimized Curcuma longa extract is efficient on knee osteoarthritis pain: a double-blind multicenter randomized placebo controlled three-arm study. *Arthritis Research & Therapy.* 2019;21(1):179. doi:10.1186/s13075-019-1960-5
34. Wang Z, Jones G, Winzenberg T, et al. Effectiveness of Curcuma longa Extract for the Treatment of Symptoms and Effusion–Synovitis of Knee Osteoarthritis. *Ann Intern Med.* 2020;173(11):861-869.



Compass Opioid Prescribing + Treatment Guidance Toolkit

doi:10.7326/M20-0990

35. Etzel R. Special extract of *BOSWELLIA serrata* (H 15) in the treatment of rheumatoid arthritis. *Phytomedicine*. 1996;3(1):91-94. doi:10.1016/S0944-7113(96)80019-5
36. Kimmatkar N, Thawani V, Hingorani L, Khiyani R. Efficacy and tolerability of *Boswellia serrata* extract in treatment of osteoarthritis of knee – A randomized double blind placebo controlled trial. *Phytomedicine*. 2003;10(1):3-7. doi:10.1078/094471103321648593
37. Prabhavathi K, Chandra USJ, Soanker R, Rani PU. A randomized, double blind, placebo controlled, cross over study to evaluate the analgesic activity of *Boswellia serrata* in healthy volunteers using mechanical pain model. *Indian J Pharmacol*. 2014;46(5):475-479. doi:10.4103/0253-7613.140570
38. Zhang Z, Leong DJ, Xu L, et al. Curcumin slows osteoarthritis progression and relieves osteoarthritis-associated pain symptoms in a post-traumatic osteoarthritis mouse model. *Arthritis Research & Therapy*. 2016;18(1):128. doi:10.1186/s13075-016-1025-y
39. Hewlings SJ, Kalman DS. Curcumin: A Review of Its Effects on Human Health. *Foods*. 2017;6(10):92. doi:10.3390/foods6100092
40. Sun J, Chen F, Braun C, et al. Role of curcumin in the management of pathological pain. *Phytomedicine*. 2018;48:129-140. doi:10.1016/j.phymed.2018.04.045
41. Rohdewald PJ. Update on the clinical pharmacology of Pycnogenol(R). *Medical Research Archives*. 2015;(3). Accessed July 30, 2021. <https://esmed.org/MRA/mra/article/view/183>
42. Rohdewald PJ. Review on Sustained Relief of Osteoarthritis Symptoms with a Proprietary Extract from Pine Bark, Pycnogenol. *Journal of Medicinal Food*. 2018;21(1):1-4. doi:10.1089/jmf.2017.0015
43. Nascimento OJM, Pessoa BL, Orsini M, et al. Neuropathic Pain Treatment: Still a Challenge. *Neurology International*. 2016;8(2):36-38. doi:10.4081/ni.2016.6322
44. Di Stefano G, Di Lionardo A, Galosi E, Truini A, Cruccu G. Acetyl-L-carnitine in painful peripheral neuropathy: a systematic review. *J Pain Res*. 2019;12:1341-1351. doi:10.2147/JPR.S190231
45. Lugo JP, Saiyed ZM, Lau FC, et al. Undenatured type II collagen (UC-II®) for joint support: a randomized, double-blind, placebo-controlled study in healthy volunteers. *Journal of the International Society of Sports Nutrition*. 2013;10(1):48. doi:10.1186/1550-2783-10-48
46. Usha PR, Naidu MUR. Randomised, Double-Blind, Parallel, Placebo-Controlled Study of Oral Glucosamine, Methylsulfonylmethane and their Combination in Osteoarthritis. *Clin Drug Investig*. 2004;24(6):353-363. doi:10.2165/00044011-200424060-00005
47. Almezgagi M, Zhang Y, Hezam K, et al. Diacerein: Recent insight into pharmacological activities and molecular pathways. *Biomedicine & Pharmacotherapy*. 2020;131:110594. doi:10.1016/j.biopha.2020.110594
48. Ruane R, Griffiths P. Glucosamine therapy compared to ibuprofen for joint pain. *Br J Community Nurs*. 2002;7(3):148-152. doi:10.12968/bjcn.2002.7.3.10214
49. Braham R, Dawson B, Goodman C. The effect of glucosamine supplementation on people experiencing regular knee pain. *British Journal of Sports Medicine*. 2003;37(1):45-49. doi:10.1136/bjsm.37.1.45



Compass Opioid Prescribing + Treatment Guidance Toolkit

50. Monfort J, Martel-Pelletier J, Pelletier J-P. Chondroitin sulphate for symptomatic osteoarthritis: critical appraisal of meta-analyses. *Current Medical Research and Opinion*. 2008;24(5):1303-1308. doi:10.1185/030079908X297231
51. Knotkova H, Pappagallo M, Szallasi A. Capsaicin (TRPV1 Agonist) Therapy for Pain Relief: Farewell or Revival? *The Clinical Journal of Pain*. 2008;24(2):142-154. doi:10.1097/AJP.0b013e318158ed9e
52. Tenreiro Pinto J, Pereira FC, Loureiro MC, Gama R, Fernandes HL. Efficacy Analysis of Capsaicin 8% Patch in Neuropathic Peripheral Pain Treatment. *Pharmacology*. 2018;101(5-6):290-297. doi:10.1159/000487444
53. Vallianou N, Evangelopoulos A, Koutalas P. Alpha-Lipoic Acid and Diabetic Neuropathy. *Rev Diabet Stud*. 2009;6(4):230-236. doi:10.1900/RDS.2009.6.230
54. McIlduff CE, Rutkove SB. Critical appraisal of the use of alpha lipoic acid (thioctic acid) in the treatment of symptomatic diabetic polyneuropathy. *Ther Clin Risk Manag*. 2011;7:377-385. doi:10.2147/TCRM.S11325
55. Shell WE, Charuvastra EH, DeWood MA, May LA, Bullias DH, Silver DS. A Double-Blind Controlled Trial of a Single Dose Naproxen and an Amino Acid Medical Food Theramine for the Treatment of Low Back Pain. *American Journal of Therapeutics*. 2012;19(2):108-114. doi:10.1097/MJT.0b013e3181f4b297
56. Ehrlich GE. Myofascial Pain and Fibromyalgia: Trigger Point Management. *JAMA*. 1995;274(4):351-352. doi:10.1001/jama.1995.03530040079049
57. Alvarez DJ, Rockwell PG. Trigger Points: Diagnosis and Management. *AFP*. 2002;65(4):653.
58. Tough EA, White AR, Cummings TM, Richards SH, Campbell JL. Acupuncture and dry needling in the management of myofascial trigger point pain: A systematic review and meta-analysis of randomised controlled trials. *European Journal of Pain*. 2009;13(1):3-10. doi:10.1016/j.ejpain.2008.02.006
59. Kietrys DM, Palombaro KM, Azzaretto E, et al. Effectiveness of Dry Needling for Upper-Quarter Myofascial Pain: A Systematic Review and Meta-analysis. *J Orthop Sports Phys Ther*. 2013;43(9):620-634. doi:10.2519/jospt.2013.4668
60. Mellick LB, McIlrath ST, Mellick GA. Treatment of Headaches in the ED With Lower Cervical Intramuscular Bupivacaine Injections: A 1-Year Retrospective Review of 417 Patients. *Headache: The Journal of Head and Face Pain*. 2006;46(9):1441-1449. doi:10.1111/j.1526-4610.2006.00586.x
61. Mellick L, Verma N. Headache Management with Occipital Nerve Blocks, Cervical Injections and Trigger Point Injections. *The Open Emergency Medicine Journal*. 2010;3(1). Accessed March 19, 2020. <https://benthamopen.com/ABSTRACT/TOEMJ-3-32>
62. Espejo-Antúnez L, Tejeda JF-H, Albornoz-Cabello M, et al. Dry needling in the management of myofascial trigger points: A systematic review of randomized controlled trials. *Complementary Therapies in Medicine*. 2017;33:46-57. doi:10.1016/j.ctim.2017.06.003
63. Johnson MI. Transcutaneous Electrical Nerve Stimulation (TENS). In: ELS. American Cancer Society; 2012. doi:10.1002/9780470015902.a0024044
64. Vance CG, Dailey DL, Rakel BA, Sluka KA. Using TENS for pain control: the state of the evidence.



Compass Opioid Prescribing + Treatment Guidance Toolkit

Pain Management. 2014;4(3):197-209. doi:10.2217/pmt.14.13

65. Gibson W, Wand BM, Meads C, Catley MJ, O'Connell NE. Transcutaneous electrical nerve stimulation (TENS) for chronic pain - an overview of Cochrane Reviews. Cochrane Database of Systematic Reviews. 2019;(4). doi:10.1002/14651858.CD011890.pub3
66. Kerai S, Saxena KN, Taneja B, Sehrawat L. Role of transcutaneous electrical nerve stimulation in post-operative analgesia. Indian J Anaesth. 2014;58(4):388-393. doi:10.4103/0019-5049.138966

This material was prepared by the Compass Healthcare Collaborative, the Opioid Prescriber Safety and Support contractor, under contract with the Centers for Medicare & Medicaid Services (CMS), an agency of the U.S. Department of Health and Human Services. Views expressed in this material do not necessarily reflect the official views or policy of CMS or HHS, and any reference to a specific product or entity herein does not constitute endorsement of that product or entity by CMS or HHS.

Developed in collaboration with Stader Opioid Consultants.