



MS Contin 30 mg + IR Opioid to Buprenorphine

How to Cross Taper from Your MS Contin 30 mg Tablets +/- Short-Acting Opioid to Buprenorphine


Are you taking MS Contin 30 mg tablets by mouth twice daily, plus one of the following opioid medications?

- + Norco (hydrocodone/acetaminophen) 10/325 mg tabs: 1 tab by mouth 3-4x daily
- + Percocet (oxycodone/acetaminophen) 10/325 mg tabs: 1 tab by mouth 3-4x daily
- + Oxycodone IR 10 mg tabs: 1 tab by mouth 3-4x daily

If you are using MS Contin and taking one of these medications, or something similar, you may be appropriate to switch to a safer, more effective pain management medication called **Suboxone**, or **buprenorphine** (+/- naloxone). You can be transitioned from your opioid to buprenorphine slowly over two weeks, working with your provider to make sure your pain is controlled while avoiding significant withdrawal.

1. You will be prescribed NALOXONE. You and your support member/s will be counseled on appropriate use of the naloxone prior to beginning the taper. Naloxone is a medication that reduces the risk of death from taking too many opioids. This is a key safety measure.
2. Learn how to use the Suboxone medication. The table below shows a visual of how to use the suboxone 2 mg – 0.5 mg SL films each day during week 1 of the cross taper:

2 – 0.5mg Suboxone Film



The first strip will be cut into 2 pieces



Half of it is then cut into 2 pieces (1/4 of a strip).



Compass Opioid Prescribing + Treatment Guidance Toolkit

		AM		PM	Date (write in)
1	¼ film	☐	-		
2	¼ film	☐	¼ film	☐	
3	½ film	☐	½ film	☐	
4	1 film	☐	1 film	☐	
5	1 ½ film	☐ ☐	1 ½ film	☐ ☐	
6	2 films	☐ ☐	2 films	☐ ☐	
7	2 – 3 films	☐ ☐ ☐	2 – 3 films	☐ ☐ ☐	

3. Plan out a reasonable **cross taper schedule**. The table below shows how to increase Suboxone and decrease the MS Contin over the first week, followed by decreasing the short-acting (ie immediate release [IR]) opioid over the second week. For representation purposes, the table will assume the patient is taking a concomitant oral opioid IR at 10 mg (+/- APAP) per dose 3-4x daily. The titration of the Suboxone during the second week and beyond will be very patient-dependent, and your provider will work closely with you to find the best regimen.

Time Point	Buprenorphine Microinduction Recommendation	
	Suboxone Rec	MS Contin + Opioid IR Rec
Day 1 (Initial Ap)	(1/4 film) SL daily	MS Contin 30 mg 2x daily; Continue opioid IR 10 mg tab: 1 tab 3-4x daily
Day 2	(1/4 film) SL 2x daily	Continue
Day 3	(1/2 film) SL 2x daily	Continue
Day 4	1 film SL 2x daily	MS Contin 30 mg 1x daily; Continue opioid IR 10 mg tab: 1 tab 3-4x daily
Day 5	1.5 film SL 2x daily	Continue
Day 6	2 films* SL 2x daily	Continue
Day 7 (F/up apt)	2 -3 films* SL 2x daily	Stop use of MS Contin; Continue opioid IR 10 mg tab: 1 tab 3-4x daily
Day 8	2-4 films* SL 2-3x daily	Reduce to opioid IR 5 mg tab**: 1.5 tabs 3-4x daily
Days 9-11	Based on craving/pain response: up to 16mg-4mg to 24mg-6mg/day split 3-4x daily	Reduce to opioid IR 5 mg tab: 1 tab 3-4x daily
Days 12-13		Reduce to opioid IR 5 mg tab: 0.5 tab 3-4x daily
Day 14 (F/up apt)		STOP oral opioid or continue as needed dosing for additional pain relief
Days 15 – beyond		

*Based on pain response; may not need to increase Suboxone dose any higher than 2 films per dose at this point, vs. increasing frequency to 3 or 4x daily. For best pain relief, 3 or 4x daily dosing is recommended. Rarely will a patient on these opioid doses need Suboxone doses of 3 or 4 films SL at a time.

**Take careful note of IR opioid tablet strength 10 mg vs 5 mg.