



Compass Opioid Prescribing + Treatment Guidance Toolkit

Oxycodone IR Taper

How to Taper Your Immediate-Release Oxycodone Medication

Are you taking the following opioid medication?

- + Percocet (oxycodone/acetaminophen) 10/325 mg tabs: 1-2 tab(s) by mouth 3-4x daily
- + Oxycodone IR 10 mg tabs: 1-2 tab(s) by mouth 3-4x daily

If you are taking one of these medications, or something similar, you may be appropriate to taper to a safer dosage or off altogether. You can be tapered down or off your oxycodone slowly, working with your provider to make sure your pain is controlled while avoiding significant withdrawal.

1. You will be prescribed NALOXONE. You and your support member/s will be counseled on appropriate use of the naloxone prior to beginning the taper. Naloxone is a medication that reduces the risk of death from taking too many opioids. This is a key safety measure.
2. Talk to your provider about other nonopioid medications and nonpharmacologic treatments that can help augment pain and mitigate withdrawal.
3. Use a chart, calendar, and/or medication box to help keep you on track with the correct dosage each week.
4. Plan out a reasonable **taper schedule**. The table below shows how to decrease your IR oxycodone product over a 1+ year period. This schedule can be accelerated based on individual response and conversations with your doctor. The goal is to get you to the finish line safely, not how quickly you get there!

Oxycodone IR Product Taper Recommendation					
Week	Dose 1	Dose 2	Dose 3	Dose 4	Total MME
0	20 mg	20 mg	20 mg	20 mg	120
1-4	20 mg	15 mg	15 mg	20 mg	105
5-8	15 mg	15 mg	15 mg	15 mg	90
9-12	15 mg	10 mg	10 mg	15 mg	75
13-16	10 mg	10 mg	10 mg	10 mg	60
17-20	7.5 mg	7.5 mg	7.5 mg	7.5 mg	45
21-24	5 mg	5 mg	5 mg	5 mg	30*
25-30	5 mg	2.5 mg	5 mg	5 mg	26.25
31-36	5 mg	2.5 mg	2.5 mg	5 mg	22.5
37-42	2.5 mg	2.5 mg	2.5 mg	5 mg	18.75
43-48	2.5 mg	2.5 mg	2.5 mg	2.5 mg	15
49-54	2.5 mg	2.5 mg		2.5 mg	11.25
54-60	2.5 mg		2.5 mg		7.5
61-62	2.5 mg				3.75

*Taper slows once reaching ~20% of total starting dose.



Compass Opioid Prescribing + Treatment Guidance Toolkit

Dosing Help

20 mg dose = 2 of the 10 mg tabs

15 mg dose = 1.5 of the 10 mg tabs

7.5 mg dose = 1.5 of the 5 mg tabs

2.5 mg dose = 0.5 of the 5 mg tabs

This material was prepared by the Compass Healthcare Collaborative, the Opioid Prescriber Safety and Support contractor, under contract with the Centers for Medicare & Medicaid Services (CMS), an agency of the U.S. Department of Health and Human Services. Views expressed in this material do not necessarily reflect the official views or policy of CMS or HHS, and any reference to a specific product or entity herein does not constitute endorsement of that product or entity by CMS or HHS.

Developed in collaboration with Stader Opioid Consultants.